

M.R.No. : _____

Date : _____

Age : _____ Sex : M / F

Weight : Kg

Anaesthesia Record

Name : _____

Operative Planned

Surgeon : Dr. _____

Anaesthetist : Dr. _____

Nurse : _____

Preoperative

H/O any prolonged illness

H/O Recent respiratory infection

H/O Previous Surgery

H/O Previous anaesthesia

H/O Allergy to drugs

H/O Diabetes / Hypertension / IHD / Asthma

Drug History

Cortisone

Antibiotic

Antihypertensive

Cardiac

Antidiabetic

Antiepileptic

Examination

Temp : _____ PR : _____ BP : _____

CVS _____ RS _____

Investigation

HB

Urine Albumin

Urine Sugar

X-Ray - Chest

ECG

Blood Urea

Blood Sugar

Others

Premedication

Time _____

Atropine Pyrolate Midazolam

Operative

PR : _____ BP : _____

Mask ETT LMA

Drugs

Thio Propofol Ketamine

Suxa Vec Attacurium

N₂O O₂ Sevo / Halo

Monitoring

Pulse Oximeter

Fentanyl Pentazocine

ECG

Metoclopramide Ondansetron

Non Invasive BP

Reversal : Glycopyrolate + Neostigmin

Incidents if any

- Difficult intubation
- Fall in O₂ saturation <90
- Rise in BP > 20%
- Fall in heart rate > 20%
- Rise (or) fall in temp
- Hypotension
- Extubation spasm
- Nausea & Vomiting after extubation

Post Operative

On Admission into recovery room

Conscious Level

- Unconscious
- Responds to deep pain
- Responds to minimal pain
- Responds to command
- Conscious and alert

Restlessness

Vomiting

SpO₂

HR

RR

Time to recover :

Chart

