


**Failure to correct refractive error leads to decreased vision and also hampers the academic activities of the child.**

**Children found with symptoms of refractive errors should be brought to an ophthalmologist for necessary correction.**



**Parents should encourage their children to wear glasses. Wearing glasses is not a stigma.**




**Loading children with vitamin A may neither help in preventing nor correcting refractive errors.**

**Spectacles perse do not increase or decrease the power.**

## Normal vision

In normal vision light rays from an object focus on the retina.



## Refractive error

Refractive error is said to exist when the light rays get focused in front or behind the retina causing blurred vision.

## Causes

- The eyeball being larger or smaller than the normal size.
- Changes in the crystalline lens.
- Flat or steeper corneal curvature.

## Types of refractive errors

The common types of refractive errors that occur in children are myopia (near sightedness), hyperopia (far sightedness) and Astigmatism.

## Hyperopia


In hyperopia or farsightedness, the light rays from an object form an image behind the retina because of the small size of the eye ball

## Myopia

The light rays from an object form an image in front of the retina. This occurs when the cornea is curved too much or if the eye is too long.


## Astigmatism

This occurs due to changes in corneal curvatures.



## Symptoms of Refractive Errors

- Children can have difficulty in reading small letters on the blackboard.
- Some children squeeze their eyes while trying to see distant objects like blackboard, television etc.
- Children hold books close to their face while reading.






- Children with myopia have defective vision for distance and clear vision for near.
- They can experience eye strain while trying to read for long hours.
- Some children with hyperopia can present with squint.
- Children can develop swellings on the lids due to constant rubbing of the eyes to see things clearly.
- Eye pain and headaches may occur.



The child should be immediately brought to an ophthalmologist, if any one of the above symptoms is observed.

### Treatment for Refractive Errors

- Correction using spectacles is the best option available for refractive errors.
- The power of the glasses may change depending on the growth of the eye ball.
- An eye checkup has to be done once in 6 months for children under 5 years of age and once a year thereafter and change of glasses when necessary.
- Children older than 15 years can use contact lenses if they don't want spectacles. Those over 18 years of age with stable power also have the option of lasik, a laser refractive procedure apart from contact lenses.
- Glasses should be worn constantly according to the advice of your ophthalmologist.



**Help your child see the world better.**

**Wearing glasses is not a stigma.....**

**Supported by**



C-10, Ground Floor,  
Gulmohar Park,  
New Delhi - 110 049,  
India  
Phone : +91 11 2696 0513  
<http://www.orbis.org>



1, Anna Nagar, Madurai-625 020, Tamil Nadu, India  
Phone : 0452-5356 100; Fax : 91-452-253 0984  
E-mail : [childreneye@aravind.org](mailto:childreneye@aravind.org)  
Website : [www.aravind.org](http://www.aravind.org)

*Refractive  
Errors  
in  
Children  
To be corrected  
or ignored?*

